

## THE GUILD OF SAWGRASS-INHABITING ANTS IN THE FLORIDA KEYS\*

BY BLAINE J. COLE  
Department of Biology,  
University of Utah,  
Salt Lake City, UT 84112

A guild is a group of species using similar resources in a similar manner (Root 1967). The guild of ants on which I report here uses, as nest sites, the hollow stems of sawgrass (*Cladium jamaicense*). The primary objectives of this study were to examine this guild for the purposes of: 1. ascertaining the extent to which there is evidence of competition for nest sites. 2. determining the extent to which available nest sites are filled. 3. determining whether *Solenopsis picta* (Emery) nests in association with other species.

### Materials and Methods

I examined 119 dead, erect culms of sawgrass, *Cladium jamaicense*, on Sugarloaf Key in Monroe County, Florida. Data were collected between 7/29/81 and 8/3/81. Each stem had seven or more internodal regions. For the ants in sawgrass stems the following data were recorded: the species present, the inside diameter of the internodal segments occupied and the internodal segment in which the colony was housed. For *Pseudomyrmex pallida* (F. Smith) the number and location of queens were also recorded. The inside diameter of a culm was measured with a micrometer to the nearest 0.1 mm. The internodal segments were numbered with the lowest segment numbered one. The numbered segments indicated relative height on the culm. Due to individual variation in the height of *C. jamaicense*, this does not translate directly into absolute height.

### Results

Out of 119 sawgrass culms examined, 34 (29%) did not have a colony of any species. The occurrences of various species as well as

---

\*Manuscript received by the editor September 10, 1982

their co-occurrences with other species is given in Table 1. *Pseudomyrmex pallida* is by far the most frequent ant, found in 57 culms or 48% of the total. *Tapinoma littorale* (Wheeler) and *Solenopsis picta* are each found in approximately 10% of the total.

The co-occurrence of series are also given in Table 1. *Solenopsis picta* is found frequently with other species including *P. pallida*, *Zacryptocerus varians* (F. Smith), and *Camponotus planatus* (Roger). No other ant species co-occur with *P. pallida*. *T. littorale* is the only species frequent enough to examine statistically. If *P. pallida* and *T. littorale* assorted into culms independently of one another, the expected number of co-occurrences would be 5.75. *T. littorale* never co-occurs with *P. pallida*, a difference that is statistically significant ( $X^2 = 11.0$ ,  $p < 0.001$ ).

It is possible to calculate the probability that *S. picta* should be the only species that co-occurs with *P. pallida*. This calculation can be done independently of our knowledge that *T. littorale* is negatively associated with *P. pallida* ( $p = 0.002$ ) or contingent on our knowledge of this relationship ( $p = 0.02$ ). In either case, it is shown that if any species occurs with *P. pallida* it is likely to be *S. picta*. It is not possible to statistically demonstrate the stronger statement that *S. picta* is positively associated with *P. pallida*. Indeed, this seems not to be the case due to co-occurrence of *S. picta* and other species.

If one combines the data of *Z. varians*, *C. planatus*, *Leptothorax allardycei* (Mann) and *Pseudomyrmex elongatus* (Mayr), one can also demonstrate that this aggregate is negatively associated with *P. pallida* ( $X^2 = 12.0$   $p < 0.001$ ). Due to the relative rarity of these species, one cannot test each species individually. This result must be considered tentative.

Table 2 gives some characteristics of the nests of the guild of sawgrass inhabiting ants. The average inside diameter and standard deviation of internodes occupied by *P. pallida* is calculated separately for that subset of the colonies that occupy a single internode and for that subset that occupy more than one internode. Nests of *P. pallida* that occupy a single internode have an inside diameter of 2.38 mm. The inside diameter of internodes occupied by *P. pallida* that are found in two internodes are 2.45 and 2.03 for the lower and upper chamber respectively. The diameter of the single nest chamber does not differ from that of the lower nest chamber of a *P. pallida* colony that occupies two chambers ( $t_s = 0.53$ ,  $p > 0.5$ ).

Table 1. Co-occurrence of Sawgrass Ants.

Species found:	no other ant	Co-occurring with:					Total
		P.p.	T.l.	S.p.	Z.v.	C.p.	
<i>Pseudomyrmex pallida</i>	52	—	0	5	0	0	57
<i>Tapinoma littorale</i>	10	0	—	0	1	0	12
<i>Solenopsis picta</i>	2	5	0	—	3	1	11
<i>Zacryptocerus varians</i>	2	0	1	3	—	0	7
<i>Camponotus planatus</i>	3	0	0	1	0	—	4
<i>Leptothorax allardycei</i>	0	0	1	0	1	0	2
<i>Pseudomyrmex elongatus</i>	1	0	0	0	0	0	1
Unidentified spider	1	1	0	0	0	0	2
Nothing	—	—	—	—	—	—	34

Table 2. Nest Characteristics of Sawgrass Ants

Species		Nest Characteristics		
		Inside Diameter mean (sdev, n)	Internode Occupied	# internodes occupied/ culm
<i>Pseudomyrmex pallida</i>	sgl. chamber	2.38 (.41, 39)	3.7	1.4
	lower	2.45 (.41, 13)	3.7	
	dbl. chamber			
	upper	2.03 (.37, 13)	4.7	
<i>Solenopsis picta</i>		3.41 (1.04, 12)	2.2	1.2
<i>Tapinoma littorale</i>		2.45 (.63, 15)	4.3	1.15
<i>Camponotus planatus</i>		4.73 (.82, 7)	2.3	—
<i>Zacryptocerus varians</i>		3.13 (.61, 8)	3.6	—

The average inside diameter is greatest for *C. planatus* (4.73), surprisingly large for *S. picta*, which is such a minute ant, and smallest for *P. pallida* (2.38, single chamber). The average diameter of internodes occupied by *P. pallida* and *T. littorale* does not differ significantly ( $t$ -test,  $t_s = .70$ ,  $p > 0.5$ ).

As shown in Table 2 the internode occupied by the nest parallels the results of internode diameter. Since larger internodes are lower

on the culm, species that inhabit internodes with large diameter also inhabit low internodes.

The number of internodes occupied per culm is given in Table 2 for *P. pallida*, *S. picta*, and *T. littorale*. *P. pallida* has a tendency to occupy more internodes per culm (1.4) than does either *S. picta* (1.2) or *T. littorale* (1.15).

In most *P. pallida* nests a queen was located. However, in 19% of the nests a queen was not seen. It is conceivable that the queen could have been overlooked in these nests. In the 46 nests in which a queen was noted, 31 (67%) had a single queen, 8 (17%) had two queens, 6 (13%) and one had four queens. When multiple queens are found in nests occupying multiple chambers, there is no tendency for the queens either to be found in a single chamber or to disperse to separate chambers. When a single queen is found in a nest occupying multiple internodes there is a tendency for the queen to occupy the higher internode.

### Discussion

The guild of sawgrass inhabiting ants is a collection of species for which there is evidence that certain pairs of species compete for nest sites and certain pairs of species do not. *P. pallida* and *T. littorale* are strongly negatively associated. This pair of species was not encountered inhabiting the same sawgrass culm. *P. pallida* and *T. littorale* occupy internodes of similar physical characteristics (inside diameter, and relative height on the culm). It is less likely that the two species compete for an internode of particular character than they compete for the space of an entire culm (Levings and Traniello 1981, Cole 1982).

*P. pallida* shows no evidence of competition for nest sites with *S. picta*. The distributions of *S. picta* and *P. pallida* are independent of one another. These two species are found in the same sawgrass culm with *S. picta* occupying larger and lower internodes. There is little evidence to suggest that *S. picta* is found in association with other species of ants. It seems to be found frequently in association with *P. pallida* simply due to the fact that *P. pallida* is common. *S. picta* has been referred to as a thief ant which nests in close proximity to other ants and specializes in stealing brood from them. In approximately half of the cases in which *S. picta* is found in a sawgrass culm

with another species of ant, there is at least one intervening, empty internode between *S. picta* and the other species.

Of the total sawgrass culms, 71% are occupied by at least one species. Let C be the average probability that a species will colonize a sawgrass culm and E be the average probability that a colony will go extinct. Then the equilibrium fraction of sawgrass culms occupied is  $C/C+E = 0.71$ . One can then obtain an estimate of the rate of extinction relative to the rate of colonization as  $C = 2.5E$ .

If colonization takes place on an annual cycle, then one can estimate that the average lifespan of a colony which becomes established is about 2.5 years. This estimate assumes that the occupancy of sawgrass culms is at equilibrium. In addition, data from several species, each of which may not have the same demographic characteristics, are combined. This is not as bad as it seems, however, due to the fact that the bulk of the species' occurrences are of *P. pallida*. The estimate of average colony longevity is principally an estimate based on *P. pallida*.

The inside diameter and position of the lower nest chamber of *P. pallida* colonies that inhabit two internodes is comparable to the inside diameter and position of the internode occupied by *P. pallida* in a single nest chamber. This suggests that *P. pallida* move up to occupy a second internode. The fact that the queen of *P. pallida* tends to be found in the upper chamber suggests that the queen moves into the newer, smaller or higher nest chamber.

The average inside diameter of sawgrass culms occupied by *Z. varians* (3.13 mm) is not significantly different from the average inside diameter of hollow stems of red mangrove occupied by *Z. varians* (2.95 mm, Cole 1979,  $n = 114$ ,  $t_s = 0.94$ ,  $p > 0.2$ ). The major workers of *Z. varians* are morphologically modified for passive colony defense (Wilson 1976, Cole 1980). It is reasonable to suppose that there is selective pressure of *Z. varians* to choose, as nest sites, hollow stems that are of a suitable size to allow the major workers to block off the stem and bulldoze out intruders.

This research supported, in part, by a grant from Sigma Xi.

#### REFERENCES

- COLE, B. J. 1979. Assembly of mangrove ant communities. Ph.D. dissertation. Princeton University. vi + 123 pp.

- COLE, B. J. 1980. Repertoire convergence in two mangrove ants, *Zacryptocerus varians* and *Camponotus (Colobopsis)* sp. *Insectes Sociaux* **27**: 265-275.
- COLE, B. J. 1982. Assembly of mangrove ant communities: patterns of geographical distribution. *J. of Anim. Ecol.* (in press).
- LEVINGS, S. C. AND J. F. A. TRANIELLO. 1981. Territoriality, nest dispersion and community structure in ants. *Psyche* **88**: 265-319.
- ROOT, R. B. 1967. The niche exploitation pattern of the blue-gray gnatcatcher. *Ecol. Monogr.* **37**: 317-350.
- WILSON, E. O. 1976. A social ethogram of the neotropical arboreal ant, *Zacryptocerus varians* (Fr. Smith) *Anim. Behav.* **24**: 354-363.